

## **SEIZURE ACTION PLAN**

Student Photo

School					
THIS STUDENT IS BEING TREATED FOR A SEIZURE SHOULD ASSIST YOU IF A SEIZURE OCCURS DURI		ION BELOW			
Student	Birthdate	Grade/Rm			
EMERGENCY CONTACTS					
Name	Relationship	Telephone number			
1	_	-			
2					
3					
Treating Physician	Tel				
Significant Medical History					
Allergies					
Triggers or warning signs					
CEIZUDE EMEDCENCY DDOTOCOL					
SEIZURE EMERGENCY PROTOCOL  A "seizure emergency" for this student is defined as:	Start Date	End Date			
☐ Seizure lasting > minutes	Start Batto	End Bute			
or more Seizures in hour(s)					
Other		<del></del>			
SEIZURE EMERGENCY PROTOC  CONTACT NURSE/CLINIC STAFF AT  Call 911 for transport to  Notify parent or emergency contact  Notify doctor  Administer emergency medications as indicated below  Other		Y AND CLARIFY BELOW)			
TREATMENT PROTOCOL DURING SCHOOL HOURS: (include daily and emergency medications)					
Daily Medication Dosage & Time of Day 0	Given Common Side I	Effects & Special Instructions			
Emergency Medication/ Instructions:					
Call 911 if  □ Seizure does not stop within m  □ Child does not start waking up within □ Child does not start waking up within □ Seizure does not stop by itself or with VNS (V.)	minutes after seizure stops minutes after seizure stops	(NO Emergency medication given) (AFTER Emergency medication is given)			
Following a seizure  ☐ Child should rest in clinic. ☐ Child may return to class (specify time frame ☐ Notify parent immediately. ☐ Send a copy of the seizure record home with child Distriction. ☐ Other		)			

**Seizure Information** - Student may experience some or all of the listed symptoms during a specific seizure.

Seiz	ure Type(s)		Description	
	Absence	•Staring •Eye blinking	•Loss of awareness •Other	
	Simple partial	•Remains conscious •Distorted sense of smell, hearing, sight	•Involuntary rhythmic jerking/twitching on one side •Other	
	Complex partial	•Confusion •Not fully responsive/unresponsive	•May appear fearful     •Purposeless, repetitive movements     •Other	
t	Generalized conic- clonic	•Convulsions •Stiffening •Breathing may be shallow •Lips or skin may have blush color	•Unconsciousness     •Confusion, weariness, or belligerence when seizure ends     •Other	
Trigg	gers or warnin	ng signs	rns to baseline in minutes.	
1		der the following circumstances		
Spec	<ul> <li>Kee</li> <li>Do</li> <li>Do</li> <li>Sta</li> <li>Rec</li> </ul> For to <ul> <li>Pro</li> <li>Kee</li> <li>Tur</li> </ul>	y calm & track time ep child safe not restrain not put anything in mouth y with child until fully conscious cord seizure in log  inic-clonic (grand mal) seizure: tect head ep airway open/watch breathing child on side  derations and Safety Precautic	A Seizure is generally considered an EMERGENCY when  A convulsive (tonic-clonic) seizure lasts longer than 5 minutes  Student has repeated seizures without regaining consciousness  Student sustains a head injury during episode  Student has a first-time seizure  Student is injured or has diabetes  Student has blue/grey color change  Student has breathing difficulties  Student has a seizure in water	
Sign	atures	Parent/Guardian Signature	 	
		i areno Guardian Signature		ty Hospitals
		Physician Signature		y Dr. Carly Wilbur 2019

Revised 2/2019 Seizures Page 2